

4. The second degree interpolating polynomial for the function $f(x) = 1/x$ on the interval $[1, 3]$ with three equally spaced nodes ($x_0 = 1, x_1 = 2, x_2 = 3$) is

$$p_2(x) = \frac{1}{6}x^2 - x + \frac{11}{6}.$$

The error formula for interpolating polynomials with equally spaced nodes is

$$|f(x) - p_n(x)| \leq \frac{h^{n+1}}{4(n+1)} \max_{\xi \in [a,b]} |f^{(n+1)}(\xi)|$$

where $h = \frac{b-a}{n}$. Use this formula to find an upper bound on the error in using the polynomial p_2 to approximate $f(x) = 1/x$ on the interval $[1, 3]$.

5. The formula for Simpson's method (not composite) on an interval is

$$\int_a^b f(x) dx = \frac{h}{3} (f(a) + 4f(m) + f(b)) - \frac{h^5}{90} f^{(4)}(\xi)$$

where $h = \frac{b-a}{2}$ and $m = \frac{a+b}{2}$ and ξ is some value between a and b . Use this rule to approximate area under the function $y = e^x$ on the interval $[0, 2]$ and estimate the error in the approximation.

6. To estimate the area under a curve using Gaussian quadrature you need to convert the function to an equivalent integral on the interval $[-1, 1]$. Then you can use Gaussian quadrature with any number of nodes. The formula for Gaussian quadrature with $n = 3$ nodes is

$$\int_{-1}^1 f(x) dx \approx \frac{5}{9}f\left(-\sqrt{\frac{3}{5}}\right) + \frac{8}{9}f(0) + \frac{5}{9}f\left(\sqrt{\frac{3}{5}}\right).$$

Find an integral on $[-1, 1]$ that is equal to $\int_0^2 e^x dx$ and then use Gaussian quadrature to estimate the value of the integral.

7. Use the centered difference quotient

$$\frac{f(x+h) - f(x-h)}{2h}$$

to approximate the derivative of $\tan x$ at $x = \pi/3$ with $h = 10^{-5}$. What is the relative error in your approximation?

8. The normal equation to find the coefficients of a (discrete) least square regression model is

$$X^T X b = X^T y.$$

Suppose you want the best fit linear function $\hat{y} = b_0 + b_1 x$ to approximate the points $(-2, 3)$, $(0, 2)$, $(2, 0)$.

- (a) What is the matrix X and the vector y in the normal equation above?

- (b) Compute $X^T X$ and $X^T y$.

- (c) Solve the normal equations to find the slope and y-intercept of the regression line $\hat{y} = b_0 + b_1 x$.

9. The Legendre polynomials are a family of orthogonal functions on the interval $[-1, 1]$. The first three Legendre polynomials are

$$P_0(x) = 1 \quad P_1(x) = x \quad P_2(x) = \frac{1}{2}(3x^2 - 1).$$

Using the Legendre functions as a basis, find the best fit (continuous least squares) 2nd degree polynomial approximation of the function $\cos x$ on the interval $[-1, 1]$. You can use the following integrals to help

$$\int_{-1}^1 P_0(x) \cos x \, dx = 1.683 \quad \int_{-1}^1 P_1(x) \cos x \, dx = 0$$

and

$$\int_{-1}^1 P_2(x) \cos x \, dx = -0.124 \quad \int_{-1}^1 P_2(x)^2 \, dx = 0.4.$$